

Research and Academics in KP California Emergency Medicine

Quarterly Report: 2026 Q2

Table of Contents

Pages

- | | |
|--|-------|
| • New Publications (n=10 + essays and abstracts) | 1-4 |
| • Just Launched or Newly Added (n=3) | 4-5 |
| • Ongoing Projects (n=42) | 6-32 |
| • Recent Publications (Jan-March 2026) | 32-33 |
-

Hot Off the Press^a

Sidney R. Garfield Exceptional Contribution Award

Dana R. Sax, MD, MPH (Oakland/Richmond) was among this year's award winners. Her research project is helping emergency physicians across all 21 medical centers rapidly diagnose and manage strokes in the eye from retinal artery occlusion.

See the 3m video: <https://tpmgawards.kaiserpermanente.org/garfield-contribution/2026/dana-sax-md-mph-2/>

Dana's publications from this project: [PubMed](#)

TPMG Publications (Northern CA)

Casey SD, Perry RM, Somers MJ, DiLena DD, Luna D, Greenhow TL, **Ballard DW**, Nguyen THP, Dalziel SR, Reed ME. [Guideline-Discordant Bronchiolitis Care in Community Emergency Departments](#). *J Pediatr*. 2026 June 16. Online ahead of print.

Grewal K, Stubblefield WB, **Casey SD**, de Wit K, **Vinson DR**, Thompson C, Hugli O, for the Emergency Advisory and Research International Board on Thrombosis (EARTH). [Preemptive anticoagulation for patients with suspected pulmonary embolism in the emergency department: An international survey of emergency physicians](#). *Acad Emerg Med*. 2026;33(4):e70297.

^a Publications, including abstracts and educational works, are organized by the region of the leading TPMG/SCPMG emergency physician author, whose name is the first one in bold font. We also highlight all KP EM co-authors. Updates for coming quarterlies can be sent to David R. Vinson, KP CREST Network: david.r.vinson@kp.org

Florio CJ, Heringer GV, Somers MJ, Qiao E, Zekar L, Middleton CE, Woldemariam ST, **Gupta N, Poth LS, Reed ME, Sperling JD, Roubinian NH, Vinson DR.** [Antenatal Pulmonary Embolism Diagnostics in Pregnant Patients with SARS-CoV-2 Infection in Community Hospitals](#). *TH Open*. 2026;10: a28721408.

Chen SJ, Sanchez AC, **Hua ST**, Murala JN, Kaushik CJ, Banga A, Dhillon GS. [Macroscopic Pulmonary Fat Embolism Secondary to Intraosseous Line Placement: A Case Report](#). *Case Rep Crit Care*. 2026:7302079.

Sax DR, Do T, Alavi M, Reed ME, Vora N, Gilbert AL, Mannis TE, Dokey AT, Vora RA. [Fundoscopic Cameras Associated with Shorter Emergency Department Length of Stay for Patients with Vision Loss](#). *Acad Emerg Med*. 2026;33(3):e70267.

Hauser KA, **Swartzberg J**, Schlessinger D, Samady N, Dove K, Donnelly M, Perkins CD, **Sax D**, Tang M, Liu VX. [Development of a predictive model for identifying high-risk older adults for geriatric emergency department screening](#). *JACEP Open*. 2026;7(4):100414.

Chen EH, **Taguma K**, Addo N, Quinn JK. [Reducing Waste: Instrument Recycling in the Emergency Department](#). *West J Emerg Med*. 2026;27(3):501-504.

Maughan BC, Jarman AF, Westafer LM, Holmes JF, Kline JA, Casey SD, Patrick N, **Vinson DR.** [Designing a Web-based Factorial Survey Experiment on Diagnostic Evaluation of Cardiovascular Symptoms in Emergency Care](#). *Res Pract Thromb Haemost*. 2026;10(4):106791.

Puebla DL, Lopez E, Kheradia T, **Zitek T**, Catapano A, Farrow RA 2nd, Kinas DH. [Accuracy of Emergency Physicians in Grading Diastolic Dysfunction Using Visual Estimation of Waveforms](#). *West J Emerg Med*. 2026;27(2):381-386.

SCPMG (Southern CA)

Seltzer J, **Moriguchi R**, Lucarelli S, Stevens CA. [Naloxone Infusions for Opioid Overdose: A 10-Year Retrospective Analysis](#). *J Emerg Med*. 2026;86:45-51.

Medical Essays

- **Meyer M.** [Shoot to Kill: The Enduring Anger of the Mass Shooter--Threat assessment can help protect healthcare workers and facilities](#). *MedPage Today*. Posted May 20, 2026.
- **Meyer M.** [What I Learned When We Treated a Possible Ebola Patient—The virus feeds off clinician fear. Preparedness and planning can help](#). *MedPage Today*. Posted May 22, 2026.
- **Meyer M.** [Five Things I Wish Someone Said at My 1998 Medical School Graduation—No one will ever care again what your class rank was](#). *MedPage Today*. Posted June 12, 2026.

Regional and National Conference Presentations

2026 Society for Academic Emergency Medicine, May 19-20, Atlanta, Georgia

Abstracts below from **CREST investigators (in blue)** are posted here: [Publications \(List\)](#) | [kpcrest](#)

1. Danielson A, Shah H, **Becht R**, McDonald R, Mumma B, Patel J. Impact of the COVID-19 Pandemic on Resident Thoracostomy Experience. *Acad Emerg Med*. 2026;33(S1):S211-S212 [abstract 487].
2. Stubblefield B, Grewat K, **Casey S**, de Wit K, **Vinson DR**, Thompson C, Hugli O. Preemptive Anticoagulation for Suspected Pulmonary Embolism: An International Survey of Emergency Physicians. *Acad Emerg Med*. 2026;33(S1):S70-S71 [abstract 147].
3. Luna D, **Casey S**, Perry R, Somers M, DiLena D, Greenhow T, **Ballard D**, Nguyen T, Dalziel S, Reed M. Guideline-discordant bronchiolitis care in Community Emergency Departments. *Acad Emerg Med*. 2026;33(S1):S287 [abstract 672].
4. **Durant E**, Warton E, **Vinson D**. Safety of Three Intravenous Rate-reducing Regimens for Atrial Fibrillation with Rapid Ventricular Response. *Acad Emerg Med*. 2026;33(S1):S89 [abstract 193].
5. Raper J, **Esener D**, Davison J, Nomura J, Boyd J, Anslip A, O'Connell N, Weekes A. Validating the Composite Pulmonary Embolism Shock Score for Predicting Clinical Deterioration. *Acad Emerg Med*. 2026;33(S1):S21 [abstract 30].
6. Picciotto N, **Feifel A**, **Viquez E**, **Lakireddy J**, Warton M, **Durant E**. Changes in Radiology Utilization After Implementation of a New Emergency Medicine Residency Program. *Acad Emerg Med*. 2026;33(S1):S212 [abstract 488].
7. **Lorenzen B**, Contreras N, Gupta A, Wei X. Racial Differences in Emergency Department Management of Patients Presenting With Chest Pain in an Integrated Health Care System. *Acad Emerg Med*. 2026;33(S1):S224 [abstract 517].
8. Mansour A, Redmond B, **Melendez A**, Auerbach M. Emergency Medicine and Neonatology Simulation Cofacilitation to Improve Neonatal Care. *Acad Emerg Med*. 2026;33(S1):S150-S151 [abstract 342].
9. Vora S, Evans L, Mansour A, **Melendez A**, Wong A, Gheasuddin A, Zhang L, Paz C, Osumah N. Virtual Reality Interprofessional Simulation to Improve Teamwork and the Culture of Safety in the Emergency Department. *Acad Emerg Med*. 2026;33(S1):S150 [abstract 339].
10. Rider A, Krzyzaniak S, Gharahbaghian L, Roszczynialski K, Karalius V, Garcia E, **Silverio L**, Ravi S. A Structured Longitudinal Curriculum to Promote Resident Engagement in Quality Improvement. *Acad Emerg Med*. 2026;33(S1):S414 [abstract 957].
11. Rider A, Daniel R, Duong D, Fee C, Krzyzaniak S, Misa N, Porter E, Roszczynialski K, Ruegner R, **Silverio L**, Sudario G. Bridging Programs, Building Leaders: The Regional Chief Resident Experience. *Acad Emerg Med*. 2026;33(S1):S413-414 [abstract 956].
12. **Vinson D**, Akkad J, Mvemba A, Wolfkind I, **Gupta N**, Rouleau S, Somers M, Pai A, Reed M, Sperling J, Roubinian N. Outpatient Care of Pregnant Patients With Acute Pulmonary Embolism in 21 U.S. Community Hospitals. *Acad Emerg Med*. 2026;33(S1):S140-S141 [abstract 317].
13. Stubblefield W, **Vinson D**, **Casey S**, Jimenez D, Bura-Riviere A, de Ancos C, García-Ortega A, del Molino F, Rivera A, Monreal M. Global Trends and Variation in Initial Anticoagulation of Patients Hospitalized With Acute Pulmonary Embolism. *Acad Emerg Med*. 2026;33(S1):S221 [abstract 509].

14. Maughan B, Westafer L, [Vinson D](#), [Casey S](#), Pulia M, Goldberg E, Baugh C, Kabrhel C, Madsen T, Peterson S, Bledsoe J, Smith A, de Wit K, Kline J, Jarman A. Implicit Bias in Diagnostic Testing for Pulmonary Embolism: The PERCEIVE-PE Study. *Acad Emerg Med*. 2026;33(S1):S232 [abstract 536].
15. Eller B, Jarman A, Kelson H, [Vinson D](#), Stubblefield W, Sebat C, Rouleau S. Intubation in patients with acute pulmonary embolism: a descriptive, retrospective cohort study. *Acad Emerg Med*. 2026;33(S1):S70 [abstract 146].
16. Campbell A, Heringer G, Smith C, Khader J, [Vinson D](#). A Cancer-Modified Pulmonary Embolism Severity Index Score Expands Outpatient Eligibility. *Acad Emerg Med*. 2026;33(S1):S142 [abstract 320].
17. Heringer G, Florio C, Campbell A, Somers M, [Vinson D](#). Antenatal Pulmonary Embolism Diagnostics in Patients With COVID-19: A Retrospective Cohort Study. *Acad Emerg Med*. 2026;33(S1):S141 [abstract 318].
18. Chen X, Farshidpour L, Wolfkind I, Gupta N, Somers M, [Vinson D](#). Prospective Validation of the Canadian Syncope Risk Score in 5 U.S. Community Hospitals: An Interim Analysis. *Acad Emerg Med*. 2026;33(S1):S89 [abstract 192].
19. Kelson H, Rouleau S, Weekes A, Kabrhel C, Mumma B, Maughan B, Eller B, Sebat C, Troha D, [Vinson D](#), Jarman A. Sex Differences Among Patients Who Received a Pulmonary Embolism Response Team Activation. *Acad Emerg Med*. 2026;33(S1):S231 [abstract 533].
20. Volz E, [Zitek T](#), Cargioli R. Mandatory vs Optional Influenza Vaccination for College Students. *Acad Emerg Med*. 2026;33(S1):S139 [abstract 314].
21. Yu A, Al-Najjar L, Hung Y, Lee-Haynes L, Mellor RG, Klein N, [Zitek T](#). Influenza-Associated Outcomes for Children in the 2024-2025 Season Stratified by Current and Previous Season Vaccination Status. *Acad Emerg Med*. 2026;33(S1):S138-139 [abstract 313].

Just Launched or Added

1. Echocardiographic Views in Rib Spaces with Differing Arm Positions

Principal Investigator: **Gabriel Rose** (San Diego)

Co-investigator: Ha-Linh Nguyen (San Diego)

Study Sites: KPSC

Summary: Using simulation models, we will assess bedside echocardiogram image quality with arm adducted vs arm abducted for various views.

Status: We have completed data collection. An abstract was presented at KP San Diego EM/FM Research Day. We submitted an abstract to the 2026 Scientific Assembly of the American College of Emergency Physicians.

2. Operator Performance of Pericardiocentesis Using Different Echocardiographic Perspectives

Principal Investigator: **Gabriel Rose** (San Diego)

Co-investigators: Ellen Chow (San Diego)

Study Sites: KPSC

Summary: Using simulation models, we will assess operator performance of pericardiocentesis from the traditional subxiphoid approach with phased array probe out of plane vs apical 4 chamber approach with the linear probe in plane. We will measure number of attempts to achieve successful pericardial puncture, time to puncture, and user survey feedback.

Status: We have completed data collection. An abstract was presented at KP San Diego EM/FM Research Day. We submitted an abstract to the 2026 Scientific Assembly of the American College of Emergency Physicians.

3. Impact of an ED triage decision support tool on accuracy of triage and timeliness of care

Principal Investigator: **Dana R. Sax** (Oakland/Richmond and the DOR)

Co-Investigators:

Funding: TPMG's Delivery Science Program

Setting: KP Northern California

Summary: This study seeks to answer this question: How is access to ED triage decision support associated with accurate patient sorting and timely care, and how can the interventions be improved prior to spread across KPNC? We will start with a pilot study in the Oakland ED to compare triage accuracy and timely care pre/post deployment. We will use a differences-in-differences approach to assess changes in outcomes at the intervention site compared with a control site/s. We will collect provider feedback on usability and workflow integration. This study has three goals: (1) Train ED nurses, physicians, and technicians on use of novel decision support tools and their integration into standard workflows; (2) Evaluate feasibility and changes in triage accuracy and timeliness of care using a pre/post design and an exploratory control site comparison; (3) Evaluate implementation effectiveness using the RE-AIM approach, including reach of the decision support; adoption of decision support recommendations and provider feedback on user experience; and impact on patient outcomes, including rates of mis-triage and delays in initial provider assignment and orders. The results will enable us to identify opportunities for iterative improvements prior to further spread.

Ongoing Research Projects^b

1. Incorporating emergency medical services (EMS) data into ED risk stratification tools: a triage-based proof of concept study

Principal Investigator: **Dustin W. Ballard** (San Rafael and Marin County EMS Director)

Co-Investigators: Mary E. Reed (DOR), Dana R. Sax (Oakland/Richmond), Greg Kann (South Sacramento and Sacramento EMS), Zita Konik (KP ED Pool Physician, Modesto, and Alameda County EMS), Tina Vitale-McDowell (KP ED Nurse Lead), and Renee Hsia (Health Services Researcher, UCSF)

Funding: TPMG Delivery Science Program (via the Physician Researcher Program)

Setting: KPNC and Marin, Oakland, and Sacramento EMS Agencies

Summary: Proper sorting of EMS patients has become increasingly important as ED patient boarding has become common. Many EMS patients wait 30 minutes or more on their gurneys before care is accepted by ED staff. This delay in ambulance patient offload time (APOT) is an urgent challenge across California: 79% of EMS offloads exceed 30 min (average 43 min). We seek to answer this question: Can EMS data from the electronic patient care record (ePCR) be incorporated with KP HealthConnect (KPHC) data to better risk stratify 911 patients in KPNC?

Aim 1: Among adults arriving via 911 EMS to KPNC EDs in 2023–2024, identify predictors—available in KPHC at the time of triage—of (1) under or over triage (defined as a mismatch between predicted and actual ED resource use) and (2) critical adverse clinical outcomes occurring within 1h of ED arrival.

Aim 2: We will integrate EMS data into the Aim 1 database for 911 EMS transports for Marin, Sacramento and Alameda counties and examine EMS predictors of Aim 1 outcomes. We hypothesize that baseline EHR data and ED triage assessment characteristics will be predictive of under or over triage and critical adverse clinical outcomes within 1h of ED arrival

Aim 3: We will use integrated data from Aims 1&2 to predict likelihood of clinical and operational outcomes of 911 EMS patients (as above).

2. Infant Fever STEWARD Project (STandardizing Emergency Work-up Around Risk Data): Three Phases

2a. Infant Fever Study (original study)

^b Active studies are organized alphabetically by the leading TPMG or SCPMG emergency physician investigator, whose name is in bold font.

Principal Investigators: **Dustin W. Ballard** (San Rafael) and Tara Greenhow (Pediatric Infectious Diseases, San Francisco)

Co-Investigators: KP CREST Network, Adam L. Sharp (DRE^c and Los Angeles), and Pediatric Hospitalists Bev Young and Tran Nguyen

Study Sites: KP Northern California and KP Southern California

Funding: Garfield Memorial Fund

Summary: We defined retrospective incidence rates of clinical and utilization outcomes in two cohorts (age 7-90 days, and 91-365 days) presenting to the emergency department (ED) in Kaiser Permanente Northern California (KPNC) and Kaiser Permanente Southern California (KPSC) with fever. We deployed these incidence data in a structured electronic clinical decision support (CDS) module that prospectively collects data. We are collecting real-time patient-specific clinical data in a structured fashion based on age strata and offer CDS links to Peds HBS/Peds ID-approved guideline documents/flowcharts. CDS content and evaluation emphasize utilization outcomes.

Status: The CDS module is in use across most KPNC facilities, used for patient care and for collecting prospective data. We have published [five manuscripts](#) total, including the project expansions noted below.

2b. California Febrile Infant Risk Stratification Tool (CA FIRST) Study

Principal Investigators: **Dustin W. Ballard** (San Rafael) and Tara Greenhow (Pediatric Infectious Diseases, San Francisco)

Co-Investigators: KP CREST Network, Beverly Young and Tran Nguyen (Pediatric Hospitalists, Roseville), Patrick Van Winkle (Pediatric Hospitalist, Anaheim), Margaret Stone (Pediatric Infectious Diseases, Woodland Hills), Sonya Negriff (Research Scientist, DRE, Pasadena)

Study Sites: KP Northern California and KP Southern California

Funding: Garfield Memorial Fund

Summary: Our CA FIRST protocol for the management of febrile infants was structured on the validated Roseville protocol and modified in light of the latest American Academy of Pediatrics (AAP) guidelines. The protocol has been accepted by physician leadership in both KP Northern and Southern CA and posted on the Clinical Library. In an earlier GMF-supported study, we designed, built, and implemented a structured electronic clinical decision support module to bring the CA FIRST protocol to EDs across KPNC. In this second phase, we are (1) retrospectively validating the CA FIRST protocol in a large cohort febrile infants in KPSC and compare its performance with the AAP guidelines; (2) prospectively evaluating the performance of the CA FIRST protocol in KPNC for key safety and diagnostic outcomes; and (3) expanding decision support in KPNC to promote best

^c DRE = KPSC Department of Research & Evaluation (Pasadena); DOR = KPNC Division of Research (Pleasanton)

practices regarding optimal ordering of chest x-rays, urinalysis and culture, and lumbar punctures in febrile infants <90 days of age.

Status: We are in analysis phase for Aims 1 and 3. We presented an abstract in May 2023 on the diagnostic yield of blood cultures in the 91-365-day population. That ms is ready for submission. Aim 2 data collection continues with over 6,400 prospective enrollments for infants aged 7-90 days.

2c. Improving clinical decision support for the care of febrile infants: expanding CA FIRST research (California Febrile Infant Risk Stratification Tool)

Principal Investigators: **Dustin W. Ballard** (San Rafael) and Tara Greenhow (Pediatric Infectious Diseases, San Francisco)

Co-investigators: From **KP Northern CA**: Dustin G. Mark (Oakland/Richmond), Mamata V. Kene (San Leandro/Fremont), David R. Vinson (Roseville/Sacramento), Tran H. P. Nguyen (Pediatric Hospital Medicine, Roseville), Madeline J. Somers, Adina S. Rauchwerger, and Mary E. Reed (DOR), with the KP CREST Network; from **KP Southern CA**: Margaret Stone (Pediatric Infectious Diseases, Woodland Hills), Joseph Colli (Pediatric Hospital Medicine, Harbor City), Sonya Negriff (DRE, Pasadena)

Study Sites: KPNC and KPSC

Funding: Garfield Memorial Fund (who has generously supported this project from its inception)

Summary: This study is expanding our ongoing work (see studies 5a/5b above) to improve the care of febrile infants with the following four aims: (1) We will retrospectively validate CA FIRST in KPNC and KPSC outpatient settings using data from 2021-2024. (2) We will implement enhanced Clinical Decision Support (CDS) in RISTRA (KPNC) with a focus on the age 91-365 day febrile infant population and the goal to decrease unnecessary utilization and treatment, specifically chest x-rays, blood cultures and antibiotics for respiratory conditions. We will prospectively track utilization and safety outcomes to assess the impact of this decision support. (3) We will determine the utility of obtaining a urine culture in the setting of negative urinalysis (UA) in infants 7-60 days. For infants 61 days to 2 years in outpatient clinics and ED, we will explore the feasibility of implementing UTICalc on the Clinical Library and embedding it into the RISTRA clinical decision support tool. We also plan to provide additional recommendations to providers on how to obtain appropriate specimens. (4) We will disseminate the data and guidelines developed in Aims 1 and 2 across KP National. The information will be shared via Clinical Library, promotion at regional chiefs' meetings, webinars and presentations with continuing medical education and social media.

Status: We have extended our investigation to the outpatient setting. We recently published risk profiles of febrile infants with COVID-19. An abstract has been submitted to Pediatric Hospital Medicine. We started prospective analysis/validation of our algorithms. We are working with KPSC on a retrospective validation. We presented two abstracts at the annual meeting of the Pediatric Academic Societies in Boston in April.

3. Non-endoscopic management of acute esophageal food impaction in the ED

Principal Investigator: Linda Lee (Gastroenterology, Sacramento)

Co-Investigators: **Sean C. Bouvet** (San Francisco) and Dan Li (Gastroenterology, Santa Clara)

Study Sites: KP Northern California

Funding: KP Northern California Community Health Research Program

Summary: Esophageal food impaction is one of the most common gastrointestinal emergencies. These patients usually require urgent endoscopy in the ED to remove the impacted food and avoid esophageal necrosis or perforation. Research regarding non-endoscopic management remains limited and the current endoscopy guidelines are over 10 years old. If non-endoscopic treatments can successfully treat patients with esophageal food impaction, it can avoid the risk, time, and expense of endoscopy. In this study, we will assess the rate of successfully resolving food impaction without endoscopy, the predictors and efficacy of non-endoscopic therapy, and the incidence of adverse events related to treatment type.

Status: We presented an abstract at Digestive Disease Week in San Diego in May. The ms is being written.

4. Predictors of Blood Culture Ordering and Results for Febrile Children in the ED

Principal Investigator: **Scott D. Casey** (Vallejo/Vacaville)

Co-investigators: Madeline J. Somers (DOR), Tara L. Greenhow (Peds ID, SFO), Mary E. Reed (DOR), Dustin Ballard (San Rafael), David R. Vinson (Roseville/Sacramento), Cole J. Florio (MS1 at California Northstate College of Medicine), Madeleine Hatch (R1 at UC Davis)

Study sites: KP Northern California

Funding: Garfield Memorial Fund

Summary: This study seeks to evaluate the characteristics associated with blood culture ordering and outcomes in febrile pediatric patients, with a specific focus on identifying predictors of true-positive results. By assessing the incidence of bacteremia and occult bacteremia in the local ED population, this research aims to fill a critical knowledge gap and provide actionable insights for clinical decision-making in the post-pneumococcal vaccine era.

Status: The manuscript is undergoing revisions for *JACEP Open*.

5. Emergency physicians' attitudes towards presumptive anticoagulation for suspected acute pulmonary embolism

Investigators: Keerat Grewal (Schwartz/Reisman Emergency Medicine Institute, Toronto, ON), **Scott D. Casey** (Vallejo/Vacaville), Olivier Hugli (Lausanne University, Switzerland), Bo Stubblefield (Vanderbilt), and David R. Vinson (Roseville/Sacramento)

Study sites: EDs across several countries

Funding: None

Summary: Consensus guidelines recommend consideration of presumptive anticoagulation in patients with a high pre-test probability of acute pulmonary embolism while pursuing diagnostic confirmation. Poor adherence to these recommendations is incompletely understood and studies assessing physicians' attitudes towards presumptive anticoagulation may illustrate barriers and facilitators to increasing presumptive anticoagulation practices through implementation science initiatives.

Status: Our first ms has been published; a second is underway: Emergency physician's risk taking, stress from uncertainty, and use of preemptive anticoagulation for suspected PE. A third paper will address preemptive anticoagulation for patients with suspected DVT.

6. Traditional ACLS training vs TeamSTEPPS plus ACLS training: a comparison of outcomes

Principal Investigator: **Charles Chiang** (San Diego)

Co-Investigators: Marlene M Alfaro, Adam Schwartz, Peter Sacchi, Sari Lahham, Ian Chong, Daniel Lee (all San Diego)

Study Site: KP San Diego

Summary: This is a prospective observational study to evaluate the effectiveness of ACLS simulation + TeamSTEPPS training compared with standard AHA didactic ACLS training received by emergency and family medicine residents. Residents that have undergone prior didactic ACLS training and are "ACLS certified" undergo a simulated cardiac arrest case in the simulation lab. Residents are then trained via simulation and TeamSTEPPS methodology on running cardiac arrest cases and are later run through another cardiac arrest simulation. Observational variables collected and compared include time to recognition of arrest, time to initiation of CPR, time to defibrillation, time to epinephrine, among others. Subjective data surveys are also collected pre- and post-simulation training.

Status: Data have been collected and analysis completed. Initial abstract completed and will be submitted for presentation at a national meeting.

7. Emergency department ultrasound-guided nerve blocks for hip fracture analgesia: a retrospective cohort study

Principal Investigator: **Edward J. Durant** (Modesto/Manteca)

Co-Investigators: Liga Yusvirazi, Kyle Herout, Nhia Yang, Valerie Lew, and Gerin River (Modesto/Manteca), Douglas Stram (DOR)

Study Sites: KP Northern California

Funding: Kaiser Permanente Northern California Graduate Medical Education, Kaiser Foundation Hospitals

Summary: This is a retrospective cohort study evaluating whether ultrasound-guided nerve blocks (UGNB) for hip fracture patients performed by emergency physicians (EPs) in the Emergency Department reduced opioid consumption compared with UGNB performed by non-EPs later in hospitalization. The study included 3,314 adult patients with hip fractures treated at KPNC hospitals between 2017 and 2022. Multivariable linear regression analysis demonstrated that EP-performed UGNB was associated with significantly lower opioid use in the first 48 hours of hospitalization, with no significant differences in complication rates between the groups.

Status: We presented an abstract at the Society for Academic Emergency Medicine in 2025. The ms is undergoing revision for the *Western Journal of Emergency Medicine*.

8. Safety of atrial fibrillation antiarrhythmic regimen using IV calcium and beta blockers

Principal Investigator: **Edward J. Durant** (Modesto and Manteca)

Co-investigators: Taylor Liu (Electrophysiology, Santa Clara), Mary Reed, Margaret Warton, and Adina Rauchwerger (KP DOR), and David R. Vinson (Roseville and Sacramento), and the KP CREST Network

Study Sites: KP Northern California

Funding: KPNC Community Health Research Program

Summary: We will determine the prevalence and outcomes in the use of combined IV rate-reducers for the acute treatment of atrial fibrillation (AF) with rapid ventricular response. Our research question is: among adult patients in the ED receiving combined therapy with IV beta blockers and calcium channel blockers for AF with rapid ventricular response, what is the 4-hour incidence of adverse events (e.g., bradycardia or hypotension requiring IV medications, cardiopulmonary resuscitation, death) compared with patients receiving monotherapy with either medication? This is a retrospective cohort study of adult patients who were treated with IV beta blockers or calcium channel blockers for a primary diagnosis of AF in the ED from 1/1/2010 to 12/31/2024. The study addresses a clear and pressing knowledge gap, as there are currently limited data on the safety of combined IV beta blockers and calcium channel blockers compared with monotherapy for rapid AF.

Status: We presented an abstract at the 2026 Society for Academic Emergency Medicine meeting in May. We are currently writing the manuscript.

9. CT Use Reduction in Ostensive Ureteral Stone (CURIOUS): retrospective validation of clinical decision rules to predict complicated ureteral stone

Principal Investigator: **Edward J. Durant** (Modesto/Manteca)

Co-Investigators: Annie Ma (UC Davis), Vignesh Arasu (Radiology, Vallejo), Raymond Bernal (Urology, Manteca), Mary E. Reed and E. Margaret Warton (DOR), Aidan Campbell (NYU), Zev Minow (Kaiser Central Valley), Cynthia Kim (California Northstate University), and David R. Vinson (Roseville/Sacramento) of the KP CREST Network.

Study Sites: KP Northern California

Funding: KP Northern California Community Health Program

Summary: Computed tomography (CT) is considered the gold standard for diagnostic imaging in suspected renal colic. Several researchers have attempted to develop clinical decision rules to predict ureteral stones without the use of CT. The main drawback of these clinical decision tools is that they were not designed to predict complications from stones, such as the need for admission or urologic intervention. In this retrospective study, we sought to derive clinical decision rules to guide imaging decisions based on the patient's risk of complicated stones. To our knowledge, ours is the first study specifically designed to derive clinical decision rules to predict clinically important stones in patients with suspected renal colic. If validated, these rules could be used to guide imaging decisions, expedite ED throughput, save resources, reduce radiation exposure, and provide a model for other EDs to follow.

Status: We published our methods paper in *Am J Emerg Med*. Our prediction rule manuscript was published in *Am J Emerg Med*: CT Use Reduction In Ostensive Ureteral Stone (CURIOUS). We are now writing a manuscript for a complementary study to evaluate how the magnitude of hydronephrosis correlates with clinically important ureteral stones. We submitted an abstract for the 2026 ACEP Research Forum meeting in October.

10. Diagnostic accuracy of retinal pathology by emergency physicians comparing ocular ultrasound and direct fundoscopic imaging

Principle Investigator: **Joshua Fuchs** (San Diego)

Co-investigators: Carlos Gonzalez-Cobos, Gabriel Rose, Dasia Esener (San Diego)

Summary: Fundoscopy can be challenging in the ED. Point-of-care ultrasonography (POCUS) offers an accurate way to diagnose emergent conditions of the eye that can replace the need for fundoscopy. Recently iCARE technology has been introduced as a more user-friendly method of performing fundoscopy. We are performing a prospective, quiz-based survey study to evaluate whether emergency physicians across a broad range of training can distinguish various pathologies of the eye based on POCUS images and iCARE fundoscopic images.

Status: Study completed. Abstract was presented at the 2024 American College of Emergency Physicians Research Forum in Las Vegas. Manuscript in preparation.

11. A retrospective analysis on the current use of bedside ultrasound in the diagnosis of acute heart failure in the emergency department

Principle Investigator: **Carlos Gonzalez-Cobos** (San Diego)

Co-investigators: Gabriel Rose, Joshua Fuchs, Dasia Esener, Peter Sacci (San Diego)

Study Site: San Diego

Summary: Congestive heart failure (CHF) is a common admitting diagnosis from the ED. Point-of-care ultrasonography (POCUS) can be highly accurate in diagnosing patients with acute CHF. POCUS for CHF using echocardiography and lung imaging is taught to residents as part of the standard core principles of training and is considered an ACGME requirement. We are performing a retrospective analysis of all CHF admissions from the ED over one year and assessing the percentage of patients who did/did not undergo POCUS as part of their diagnostic assessment.

Status: Study completed. Abstract was presented at the 2024 American College of Emergency Physicians Research Forum in Las Vegas. Manuscript in preparation.

12. ED management of alcohol withdrawal: phenobarbital vs benzodiazepines and risk of 72-hour return to the ED

Principal Investigator: **Mamata V. Kene** (San Leandro and Fremont)

Co-Investigators: Dustin G. Mark (Oakland and Richmond) and Mary E. Reed (Division of Research)

Funding: Community Health Research Program

Setting: KP Northern California

Summary: With the increasing prevalence of alcohol-use related emergency department visits, understanding the role of phenobarbital in management of mild to moderate alcohol withdrawal is important. An alternative to benzodiazepines, the traditional first line of treatment, phenobarbital has a smoother, longer metabolism that may make it better tolerated among patients who are discharged home. Prior studies of phenobarbital vs benzodiazepines either focus on inpatient endpoints or are limited in size. We propose a retrospective cross-sectional study to evaluate phenobarbital vs benzodiazepine use in the ED for alcohol withdrawal among adult ED patients with eligible alcohol withdrawal-related diagnoses AND medication administration, who are discharged home, with the primary outcome of 72 hour return to the ED. We will adjust for patient and facility characteristics that may be associated with choice of medication and severity of alcohol withdrawal. We will also evaluate the frequency of 10-day follow up in addiction medicine outpatient services among KPNC members in our sample, comparing follow up rates between patients who received phenobarbital and benzodiazepines during the index encounter. Our findings could inform the development of guidance supporting expanded phenobarbital use in the ED, including order set development and best practice guidelines.

Status: Data collection is underway.

13. Pediatric respiratory illnesses in the post-COVID era: epidemiology, ED care, and outcomes

Principal Investigator: **Mamata V. Kene** (San Leandro/Fremont)

Co-investigators: Dana R. Sax (Oakland/Richmond), Tara L. Greenhow (Pediatric ID, San Francisco), Mary E. Reed, E. Margaret Warton, and Adina S. Rauchwerger (Division of Research, Pleasanton), and the KP CREST Network

Study Sites: KP Northern California

Funding: KP Northern California Community Health Program

Summary: Pediatric respiratory illnesses are a common reason for U.S. ED visits. The COVID-19 pandemic shifted the incidence and epidemiology of respiratory diseases among children, especially in the last two years, when non-COVID respiratory infections rebounded from a pandemic dip. The rebound-associated surge tested hospital and ED capacity both at community and children's hospitals. The burden on health systems and hospitals of this changing epidemiology is not well understood. This retrospective study will evaluate pediatric respiratory-illness-related ED visits and outcomes from the pre-pandemic to current year. We seek to understand how rates of serious illness may have changed, and to identify predictors of serious illness. Our primary outcome is hospitalization; secondary outcomes include intensive respiratory support and ICU admission. We will also evaluate facility- and patient-level factors associated with serious illness and patterns of health care utilization prior to ED visits for respiratory illness. Results will inform operational planning and patient outreach and educational efforts.

Status: Our first manuscript was recently published. We are now working on our second manuscript.

14. ED note quality before and after implementation of an AI ambient listening scribe application.

Principal Investigator: **William Krauss** (San Diego)

Co-investigators: Andrew Tabone, Chelsea Tafoya, Brent Lorenzen (San Diego)

Study Site: San Diego

Summary: SCMG implemented an AI ambient listening scribe application in ambulatory and ED settings in 2024. Little is known about the effect on note quality. This study seeks to use a validated instrument (PDQI 9) to assess the quality of notes associated with ED encounters before and after adoption of the AI application.

Status: Analysts are currently extracting 580 notes that will be evaluated by our physician team to evaluate the quality of AI generated vs non-AI generated notes.

15. Pediatric Dose Optimization for Seizures in Emergency Medical Services (PediDOSE)

Principal Investigator: Manish Shah, Stanford University

Co-investigators: Multiple, including the Medical Directors of Sacramento County EMS: **Kevin E. Mackey** (TPMG emergency physician at KP South Sacramento) and long-time CREST collaborator Daniel Nishijima (Director of EM Research at UC Davis).

Funding: National Institute of Neurological Disorders and Stroke

Study Sites: 35 EMS agencies in 20 cities across the United States, including three California counties: Sacramento, Alameda and Los Angeles.

Summary: One-third of children with active seizures on ambulances arrive at EDs are still seizing. Prior research suggests that seizures managed by paramedics continue due to underdosing and delayed medication delivery. Underdosing happens when calculation errors occur, and protocols do not have evidence-based doses (~0.2 mg/kg midazolam given intranasal (IN) or intramuscular (IM)). Delayed medication delivery occurs due to the time required for dose calculation and placement of an IV line to give the medication. In this stepped-wedge, cluster-randomized trial of children aged 6m-13y with a paramedic-witnessed seizure, investigators are simplifying dosing by making it age-based, comparing effectiveness and safety of the standardized treatment protocol (intervention) vs usual care (control). If this study demonstrates that standardized, age-based IN/IM midazolam dosing is equally safe and more effective compared with current practice, the results could transform pediatric seizure treatment in the prehospital setting. More details: <https://clinicaltrials.gov/study/NCT05121324>

Status: Enrollment will continue through July.

16. Provider-level variation in head computed tomography use in ED patients with headache and misdiagnosis of critical neurologic disease.

Principal Investigator: **Dustin G. Mark** (Oakland and Richmond)

Co-Investigator: Mary E. Reed (Division of Research)

Funding: Community Health Research Program

Setting: KP Northern California

Summary: The use of neuroimaging during emergency department (ED) visits has been increasing over the past several decades. Additionally, there is wide variation in clinical practice among emergency physicians, with up to three-fold differences between providers in per-visit use of head CT. Guidelines advocate against overuse of head CT for patients with uncomplicated headache. However, it is not clear whether varying thresholds to use of head CT imaging for headache evaluations meaningfully impacts the risk of missing critical neurologic diagnoses. This knowledge gap creates a degree of uncertainty that has been cited as a principal barrier to reducing head CT utilization. To help address this knowledge gap, we propose to leverage this well-described emergency physician-level variation in the use of head CT to examine whether, among ED patients presenting with headache, exposure to an emergency physician with a higher propensity to use

head CT imaging is associated with a decreased risk of missed or delayed diagnoses of common critical neurologic diagnoses in clinical practice (ischemic stroke, subarachnoid hemorrhage, intracerebral hemorrhage, or brain tumor). Secondly, we will assess whether these varying propensities to use head CT are associated with changes in other utilization metrics (e.g. repeat ED visits). The study will be a data-only retrospective and prospective cohort study of patients aged 18 or older presenting to a KPNC ED with a chief complaint of atraumatic headache between January 1, 2016, and December 31, 2025. We will employ a preference-based instrumental variable approach to study the association between varying emergency physician-level propensity to use of head CT for headache investigation and the rate of index versus delayed diagnoses of critical neurologic pathology. This work can help refine internal efforts to address CT overutilization within the ED, which have been previously limited by a lack of covariate balancing and outcome correlation.

Status: Data collection is underway.

17. Creation of a high-fidelity 3D simulation model for performance of POCUS guided lumbar puncture and erector spinae block

Principal investigator: **Gabriel Rose** (San Diego)

Co-investigators: Dasia Esener, Drew Silver, and Eric Abrams (San Diego)

Study Site: KP San Diego

Summary: We plan to create a semi 3D printed back (lumbar) and create a ballistics gelatin in-molding to simulate the vertebrae and erector spinae muscles. We will have residents perform an LP as well as the erector spinae block under ultrasound guidance and measure pre- and post-procedural confidence.

Status: We are creating the simulation model.

18. Ultrasound guided transgluteal sciatic nerve hydrodissection for the treatment of sciatica in the emergency department: a case control cohort study.

Principal Investigators: **Gabriel Rose** (San Diego).

Co-investigators: Dasia Esener, Sara Amen (San Diego)

Summary: This study will evaluate the efficacy of using ultrasound-guided transgluteal sciatic nerve hydrodissection (TGSNH) to treat ED patients with a chief complaint of sciatica pain. Traditional practice patterns of emergency physicians (EP) to treat sciatica include a combination of NSAIDs, opioids, topical analgesics, and acetaminophen. Given the concerns related to opioid use in certain populations and the efficacy of acute relief, these treatments can provide, there is a need to investigate further acute treatments for sciatica pain. Ultrasound-guided regional anesthesia is within the scope of practice of EPs for the treatment of musculoskeletal pain. Specifically, ultrasound-guided transgluteal sciatic nerve block has been shown as an effective

treatment of sciatica in the ED; however, safety concerns regarding the potential for falls due to motor blockade or local anesthetic systemic toxicity (LAST) due to absorption of local anesthetic cannot be overlooked. Nerve hydrodissection is a technique in which a solution (e.g. D5W) is injected circumferentially around a nerve to separate it from surrounding connective tissue, which can effectively help to alleviate pain resulting from compressive neuropathy. Unlike nerve blockade with local anesthetics, nerve hydrodissection lacks the aforementioned risks of falls and LAST. Hydrodissection has been shown in numerous studies to be an effective treatment in conditions such as carpal tunnel syndrome, cervical radiculopathy, cervicogenic headache, and more recently in a case series describing its use in sciatica. Large-scale prospective studies on this technique for the treatment of sciatica in the ED are still lacking. Here we set out to study whether TGSNH is an effective and safe method to treat patients who present with sciatica to the ED.

Status: Abstract presented at 2025 ACEP Scientific Assembly, Salt Lake City, UT. Manuscript in preparation.

19. Emergency department triage support tool build and pilot

Principal Investigator: **Dana R. Sax** (Oakland/Richmond)

Co-investigators: Dustin G. Mark (Oakland/Richmond), Mary E. Reed, E. Margaret Warton, and Adina S. Rauchwerger (KP Division of Research, Pleasanton), Robert Norris (KPNC regional physician lead for ED workflow in KP HealthConnect), Carolina van Dalen (clinical informatics nurse), Tina Vitale-McDowell (Oakland RN), and the Nursing Stakeholder Advisory Team from Oakland.

Funding: Lokahi Risk Reduction Funding Initiative

Study Sites: KPNC

Summary: Over 1.2 million patients are treated annually across 21 KPNC EDs. ED nurses perform brief triage assessments that determine when patients are seen, impact who will care for the patient, and may bias diagnostic work ups. Increasing ED volumes, crowded waiting rooms, and the need to make quick decisions with limited information present significant challenges to safe and accurate triage. In our prior research, we have identified several limitations with the current triage system, including frequent mis-triage, under-recognition of the sickest patients, and meaningful delays in appropriate diagnosis and treatment due to mis-triage for patients with time-sensitive diagnoses. We propose to develop, build, and pilot test novel clinical decision support tools to support triage nurses with accurate patient sorting and early rooming of the sickest patients. We hypothesize that deployment will decrease rates of mis-triage and facilitate more timely rooming and initial diagnostic orders among critically ill patients.

- **Aim 1:** Collaborate with KPNC information technology leads and clinical stakeholders to build two new clinical decision support tools: a) Triage Snapshot tool: Brief clinical summary and patient risk profile to support triage nurses with early identification of higher risk patients; b) Waiting room prioritization tool: Real-time display of patient risk of hospitalization at triage to help teams decide which patients to room first.

- **Aim 2:** Train all KP Oakland ED triage nurses on use of tools as part of standard triage workflows.
- **Aim 3:** Perform pilot study of tools at KP Oakland. Assess feasibility of use, nurse adoption and acceptance, accuracy of triage decisions, and timeliness of care for sickest patients. Review all cases of significant under-triage.

20. Safety and operational impacts of ED mis-triage

Principal Investigator: **Dana R. Sax** (Oakland/Richmond)

Co-investigators: Dustin G. Mark (Oakland/Richmond), Mary E. Reed, E. Margaret Warton, and Adina S. Rauchwerger (KP Division of Research, Pleasanton), and the KP CREST Network

Study Sites: KP Northern California

Funding: KP Northern California Community Health Program

Summary: ED triage systems exist to sort patients based on acuity and expected resource use. Our study team previously developed and validated novel measures to identify mis-triage for patients triaged using the Emergency Severity Index. Mis-triage can be under-triage (under-recognition of disease severity or resource needs) or over-triage. Using these measures, we estimate that one-third of KPNC adult ED patients and nearly two-thirds of KPNC pediatric ED patients were mis-triaged across KPNC from 2016-2020. In this current proposal, we will build on this work to broadly characterize the potential safety, quality, and operational implications of under- and over-triage among adult and pediatric patients. Specifically, we will conduct adjusted analyses to study the association of under-triage with delays in care and downstream patient outcomes (safety and quality outcomes) and the association of over-triage with ED length of stay and delays in care (operational outcomes). We plan to conduct a sub-analysis of the impact of under-triage specifically among patients with an ED diagnosis of severe sepsis or septic shock, for whom delays in care are associated with worse patient outcomes.

Status: We have 5 publications from this study: (1) "ED triage associated with timeliness of care for patients with septic shock" was published in *Am J Emerg Med*; (2) "Association of emergency department triage accuracy on timeliness of care and patient outcomes" was published in *Ann Emerg Med*; (3) A response to a letter to the editor on the second paper above is in press with *Ann Emerg Med*; (4) "Association between pediatric ED triage accuracy and timeliness of care and patient outcomes" is in press with *Hospital Pediatrics*; (5) A final paper is in the works on triage and timely care for pediatric patients with severe sepsis/septic shock.

21. Improving safety and quality of emergency care using machine learning-based clinical decision support at triage

Principal Investigator: **Dana R. Sax** (Oakland/Richmond)

Co-investigators: Dustin W. Ballard (San Rafael), Mamata V. Kene (San Leandro/Fremont), Dustin G. Mark (Oakland/Richmond), Mary E. Reed and E. Margaret Warton (KP Division of Research, Pleasanton), and the KP CREST Network

Study Sites: KP Northern California

Funding: Agency for Healthcare Quality and Research

Summary: To manage patient visits every year, U.S. EDs prioritize allocation of limited health care resources to patients in greatest need. Most EDs use a flawed, subjective triage system which leads to mis-triage in up to one-third of patient encounters, worsening ED crowding and contributing to delays and disparities in care. ED triage research has been largely limited, having focused on hospital admission outcomes (even though 80-90% of ED visits do not result in hospital admission), and having not included pediatric visits (which represent 1 in 4 ED visits), health equity in design or prediction model evaluation, a user-centered design, or key patient safety and quality measures. This study will address these unmet needs by creating and testing a novel digital health solution using advanced predictive analytics and patient EHR data to better prioritize patient needs and acuity in the ED setting. First, we will refine triage models that predict critical illness, hospital admission, and fast-track eligibility; then measure algorithm biases and explore strategies to improve equity in triage model predictions. Second, we will map probability thresholds for each outcome into clinically relevant triage category recommendations. Using a human factors framework, we will design, build, and evaluate clinician-facing triage clinical decision support (CDS) and build the models and CDS into our EHR to efficiently display triage recommendations as part of standard workflows. Third, in a pragmatic trial across 21 hospital-based EDs and one free-standing ED, we will assess the real-time impact of CDS, measuring timeliness of care for critically ill patients, appropriate early identification of fast-track eligible patients, and ED length of stay. In addition, to test equity-driven model calibrations, we will assess bias by race, gender, and socioeconomic status. Overall, we aim to demonstrate how advanced predictive analytics and an effective user interface can be utilized at the point of care to improve ED triage by accurately predicting acuity and care complexity and prioritizing equity. Ultimately, our research will limit crowding, streamline operational flow, mitigate disparities, and lead to safer, higher quality care and better outcomes in the ED setting.

Status: Undertaking data collection

Heart Failure Research

22. KP-specific heart failure risk prediction: KPNC Standardizing Emergency Work-ups Around Risk Data (STEWARD) heart failure project

Principal Investigator: **Dana R. Sax** (Oakland/Richmond) and Mary E. Reed (DOR)

Co-investigators: Dustin G. Mark (Oakland/Richmond), Jamal Rana (Oakland), Mamata V. Kene (San Leandro/Fremont), David R. Vinson (Roseville/Sacramento), Dustin W. Ballard (San Rafael), and the KP CREST Network, with collaborators from Vanderbilt.

Study Sites: KP Northern California

Funding: The Permanente Medical Group Delivery Science Research initiative

Summary: There are over one million ED visits across the U.S. each year for acute heart failure (AHF), with an average admission rate of 84%. EDs play a major role in the care of AHF patients through symptom management, coordination of care, and risk stratification to identify sicker patients needing admission. We recently developed a clinical decision support tool to help predict AHF disease severity and in collaboration with Vinnie Liu and the Hospital Advanced Analytics Team and the HealthConnect team; the tool is now built into our EHR and ready to provide real-time risk estimates and clinical decision support. We also collected qualitative data through interviews and surveys with frontline ED providers, IT leadership, and operational and clinical leads to identify barriers and opportunities for implementation of the risk tool. We worked with cardiology and hospital-based specialist leads to develop care pathways based on patient risk.

We ran a pilot study of the risk tool with decision support in two EDs (Oakland and Richmond) to assess feasibility of extracting HF-relevant data and efficiently presenting these to ED clinicians, assess provider uptake of the tool, and assess safety of patients identified as low risk and discharged home. We collected feedback from clinicians via interviews and surveys, and updated the tool as needed prior to regional implementation and prospective evaluation.

Status: We have published [several papers](#) from this line of inquiry. STRIDE-HF has been deployed across all 21 EDs. We are now evaluating the impact of the risk tool on patient outcomes and utilization.

23. Evaluating diuresis strategies for ED patients with acute heart failure

Principal Investigator: **Dana R. Sax** (Oakland/Richmond)

Co-investigators: Dustin G. Mark (Oakland and Richmond), Jamal Rana (Cardiology, Oakland), Mary Reed and Jie Huang (DOR), and the KP CREST Network

Study Sites: KP Northern California

Funding: KPNC Community Health Research Program

Summary: We seek to understand current ED practice in diuresis of patients with acute heart failure (AHF), and to evaluate how different diuretic dosing strategies, after adjusting for other factors, are associated with patient outcomes. We plan a retrospective cohort study of all ED encounters by KPNC members with AHF who received an IV diuretic from January 1, 2023 – December 31, 2024. Patients will be identified using the natural language processing algorithm our study team previously developed and validated to improve case identification of ED patients with AHF. In Aim 1, we will use descriptive statistics to evaluate current practice around timing, dosing, and monitoring for ED diuresis; in Aim 2, we will use a multivariate logistic regression model (high versus non-high initial diuretic dose) to assess whether any patient or visit characteristics are significantly associated with initial dosing selection; in Aim 3, we will compare otherwise similar patients who receive different initial diuretic doses to assess how the dose selection is associated with patient outcomes using inverse probability weighting. Information

from this study could help inform protocols for ED diuresis, as well as strategies for effective decongestion in alternate care spaces.

Status: The manuscript is in press at *JACC Heart Fail*: “Higher emergency department diuretic dosing for acute heart failure associated with lower risk of hospitalization”

24. Improving Care by Defining the Role of ElectroNic HealTh Record-Based Alerts In a Fully Integrated Health Care Delivery SYstem for Worsening Heart Failure (IDENTIFY-WHF)

Principal Investigators: Andrew P. Ambrosy (Cardiology, San Francisco) and Alan S. Go (Division of Research)

Co-investigators: **Dana R. Sax** (Oakland/Richmond), Justin J. Slade, Van N. Selby, Jana Svetlichnaya, Ankeet S. Bhatt, and Edward J. McNulty (Cardiology, San Francisco), Amir W. Axelrod (Cardiology, Vallejo), Sirtaz Adatya and Keane K. Lee (Cardiology, Santa Clara), Harshith R. Avula (Cardiology, Walnut Creek), and Howard H. Dinh (Cardiology, South Sacramento)

KP Study Sites: KP Northern California

Funding: The Permanente Medical Group Delivery Science Research initiative

Summary: IDENTIFY-WHF is a prospective, virtual, parallel-group, randomized, quality improvement intervention of EHR-based alerts for WHF to improve the adoption of goal-directed medical treatments (GDMTs) within KPNC. Approximately 1,000 participants will be identified at KP San Francisco and Santa Clara Medical Centers during an index hospitalization or ED visit for WHF and randomized 1:1 to the intervention or usual care, stratified by service area and left ventricular ejection fraction (LVEF) category. Best practice alerts will be sent to the treating outpatient provider(s) and include (1) the most recent LVEF, vital signs, and laboratory values, (2) current GDMT, and (3) eligible GDMT. Eligible GDMT will be based on regional standards and national HF guidelines and will include the Class (Strength) of and Level (Quality) of Evidence. The primary outcome is the proportion of adults experiencing WHF with an increase in the number of prescribed GDMT classes at 30 days post-discharge from the hospital or ED.

Status: We are collecting data.

25. Non-hospital treatment venues for acute management of patients with heart failure

Principal Investigator: **Dana R. Sax** (Oakland/Richmond)

Co-investigators: Dustin G. Mark (Oakland/Richmond), Jie Huang and Mary E. Reed (DOR), and Jamal Rana (Cardiology, Oakland/Richmond)

Funding: Garfield Memorial Fund

Study Sites: KPNC

Summary: Our primary goals are to: 1) describe patients and outcomes for six common conditions (heart failure, cellulitis, pneumonia, UTI, severe anemia, and COVID-19) treated in these care spaces: observation unit/clinical decision area, ambulatory treatment centers, Advanced Care at Home, and traditional inpatient hospitalization; 2) compare risk-adjusted patient outcomes specifically for patients with acute heart failure treated in these four venues, including 30-day serious adverse events, 30-day all-cause mortality, and guideline-concordant care; and 3) compare risk-adjusted acute care utilization for patients with acute heart failure treated in these four venues, including 30-day hospital readmissions and 30-day repeat ED visits.

Status: First manuscript in preparation, data collection/analysis ongoing.

26. Impact of fundoscopic cameras in the ED on diagnostic efficiency and accuracy for patients with vision loss

Principal Investigator: **Dana R. Sax** (Oakland/Richmond), Mary Reed (DOR), and Robin Vora (Ophthalmology, Oakland/Richmond)

Co-investigators: Mubarika Alavi (DOR)

Study sites: KP Northern California

Funding: DARE's Rapid Analytics Unit Program

Summary: Among adult (age ≥ 40 years) patients who were seen in KPNC EDs with an eligible vision-related chief complaint, this study evaluated the impact of deployment of fundoscopic cameras on key process and outcome measures, including ED length of stay, need for in-person ophthalmology consultation, need for 72-hour ophthalmology follow up, and timely diagnosis of high-risk ophthalmologic conditions.

Status: We have published three studies: (1) Branch Retinal Artery Occlusion Visual Outcomes: A Community-Based Study in *Ophthalmol Retina*; (2) Branch retinal artery occlusion time to presentation and diagnosis: a retrospective review in *Int J Retina Vitreous*; (3) Fundoscopic Cameras Associated with Shorter ED Length of Stay for Patients with Vision Loss in *Acad Emerg Med*. A fourth paper is undergoing peer review.

27. To walk-in or not to walk-in: Understanding mode of arrival choices among acute stroke patients in KPNC and their effects on stroke care processes and outcomes

Principal Investigator: Mai N. Nguyen-Huynh (Neurology, Walnut Creek)

Co-PI: **Dana R. Sax** (Oakland/Richmond), Molly Burnett, Patricia Zrelak, Navdeep Sangha

Study Sites: KPNC and KPSC

Funding: Garfield Memorial Fund

Summary: There is less understanding about why patients with acute stroke symptoms choose to arrive at an ED via “walk-in” instead of via ambulance or emergency medical services (EMS), or what the effects of that decision on stroke care processes and outcomes. Aim 1 is to assess predictors of ED mode of arrival among stroke patients. Aim 2 is to determine the effects of ED arrival mode on acute stroke alert processes including thrombolytic treatment time and outcomes at discharge and at 90-day. Aim 3 is to survey patients’ knowledge about stroke risk factors, signs and symptoms, factors influencing their choice of ED arrival mode, and identify areas with opportunity for improvement.

Status: Data collection and analysis are ongoing. We presented two abstracts at the 2026 International Stroke Conference in LA in February.

28. Improving risk stratification of patients with advanced stage colorectal cancer

Principal Investigator: Faisal Cheema (Santa Clara)

Co-investigators: **Dana R. Sax** (Oakland/Richmond), Zahra Samiezade-Yazd (DOR), Janet Alexander (DOR), and Margaret Krackeler (San Francisco)

Study Sites: KP Northern California

Funding: KPNC Community Health Research Program

Summary: Our primary goals are to: 1) Describe patients recently diagnosed with advanced stage colorectal cancer receiving chemotherapy who are at high risk for a serious adverse outcome to identify patient-specific modifiable risk factors; 2) to validate a previously developed risk stratification score to predict patients at high risk of a serious adverse event; and 3) to explore improving prediction models using additional electronically available variables.

Status: Data collection is underway.

Pulmonary Embolism Research

29. Trends in initial anticoagulation of acute pulmonary embolism during hospitalization 2014-2024: Report from the RIETE registry

Principal Investigators: **David R. Vinson** (Roseville/Sacramento), Manuel Monreal (Universidad Catholica de Murcia, Murcia, Spain) and David Jimenez (Ramon and Cajal University Hospital, Madrid)

Co-investigators: William Bo Stubblefield (Vanderbilt), **Scott D. Casey** (Vallejo/Vacaville), and Manuel Monreal (Universidad Católica San Antonio de Murcia)

Study Sites: 180 sites from 31 countries across Europe, North and Latin America, and Asia.

Summary: RIETE is an ongoing, prospective multinational observational study of patients with objectively confirmed acute venous thromboembolism (VTE). The registry was begun in Spain in

2001 with the goal of gathering a large sample of patients with VTE, with specific attention to those excluded from typical randomized trials of anticoagulant therapy. This study addressed the selection of initial anticoagulation for hospitalized patients, anticoagulation trends over time, and variation between countries. Findings suggest that some regions, including the United States, could improve patient care by aligning anticoagulation practices with clinical guidelines.

Status: We presented an abstract at the Society for Academic Emergency Medicine meeting in May and will do so, as well, at the International Society on Thrombosis and Haemostasis in July. The manuscript is undergoing peer review.

30. Outpatient management of pregnancy-associated pulmonary embolism

Principal Investigator: **David R. Vinson** (Roseville/Sacramento)

Co-investigators: Madeline J. Somers and Mary E. Reed (DOR), Nareg Roubinian (Pulmonology and Critical Care, Oakland), Jeffrey D. Sperling (Maternal and Fetal Medicine, Modesto), Ashok P. Pai (Hematology/anticoagulation, Oakland), Jemma Akkad and Sam Rouleau (UC Davis resident and fellow, respectively), Audrey Mvemba (KP Oakland OB/GYN resident), and the KP CREST Network

Study Sites: KP Northern California

Funding: KPNC Community Health Research Program

Summary: Emergency physicians and obstetricians across KPNC discharge home select pregnant and postpartum patients with acute pulmonary embolism (PE) within 24 hours of their arrival to the ED and Labor and Delivery, respectively. This practice of rapid diagnostics, brief observation, and expedited discharge with close follow-up is known as outpatient management. Strangely, such a practice, though recommended by professional society guidelines, has never been described in the literature. With a 14-year retrospective cohort study across 21 medical centers from 2011 through 2024, our established multidisciplinary team (emergency medicine, maternal-fetal medicine, pulmonology, and hematology) will address 3 unexplored questions: (1) What is the prevalence of outpatient management of pregnancy-associated PE across KPNC? (2) What patient- and facility-level variables characterize those selected for outpatient care? (3) How safe is the practice as measured by the 7-day incidence of PE-related hospitalization and all-cause mortality? Results of this first-of-its-kind study will inform quality improvement measures within KPNC recommendations by professional societies.

Status: Data collection is nearing completion. We published a small case series as a teaser: Expanding outpatient management of low-risk pulmonary embolism to the pregnant population. *Eur Heart J Case Rep*. The ms on pregnant patients is under peer review. The postpartum analysis is underway.

31. Predictors of early hospital discharge in patients admitted with pulmonary embolism

Principal Investigator: Nareg H. Roubinian (Pulmonology/Critical Care, Oakland/Richmond)

Co-investigators: **David R. Vinson** (Roseville/Sacramento), Mahesh J. Balasubramanian (Adult Hospital Medicine, Roseville), Mary E. Reed, Madeline J. Somers, and Adina S. Rauchwerger (KP Division of Research, Pleasanton), and the KP CREST Network

Study Sites: KP Northern California

Funding: KP Northern California Community Health Program

Summary: Increasing outpatient management of stable ambulatory clinic and ED patients has changed the population of those being hospitalized with acute pulmonary embolism (PE). With fewer low-risk patients requiring hospitalization, patients admitted are now more complex, with higher severity of illness and more serious comorbidities. This proposal will assess the contemporary characteristics of patients requiring hospitalization for acute PE (Aim 1) and identify characteristics associated with early hospital discharge (<48h) (Aim 2). In Aim 1, we will examine characteristics of hospitalized patients admitted with PE, including trends in vital signs, oxygen requirements, and diagnostic testing at admission and during hospitalization. In Aim 2, we will examine predictors of early hospital discharge vs. prolonged hospitalization at admission and on a daily basis as part of a multivariable model. These results may inform resource allocation to support early hospital discharge or even outpatient management of patients with acute PE in the future. We hope to identify processes of care associated with safe, early discharge (e.g., early discharge on home oxygen for mildly hypoxic patients) that could be streamlined in patients who experienced prolonged hospitalization to facilitate more timely discharge.

Status: We presented an abstract at the CHEST annual meeting in October in Chicago. The manuscript is in construction.

32. Physician perceptions of risk of pulmonary embolism diagnosis using case vignettes

Principal investigator: Brandon Maughan (OHSU)

Co-investigators and study sites include, among others, Angela Jarman (UC Davis), Lauren Westafer (U Mass Baystate), Mike Pulia (UW Madison), Elizabeth Goldberg (U Colorado), Chris Baugh (Brigham), Chris Kabrhel (Mass General), Tracy Madsen (Brown), Susan Peterson (Johns Hopkins), Lauren Stewart (Indiana), and **David R. Vinson** (Roseville/Sacramento)

Summary: We will assess physician perceptions of pulmonary embolism (PE) risk using a randomized factorial experiment with a set of hypothetical patient cases with varying levels of risk for PE. We hope to (1) identify the influence of gender and racial bias in ED physicians' perceptions of PE risk and decisions to pursue diagnostic testing and (2) gather pilot data on how physicians perceive the PE risk associated with different elements of ED patient presentations (e.g., symptoms characteristics; medical history; vitals and exam). Ultimately, this study will provide pilot data for a larger study on improving PE diagnosis.

Status: We have included 650 participants across the US and Canada who have evaluated 3,000 hypothetical vignettes. The methods paper was recently published. We presented an abstract at

the Society for Academic Emergency Medicine meeting in May: Implicit Bias in Diagnostic Testing for PE. We will soon write up our findings on how patient sex affects PE diagnostics. Other mss will follow.

33. Is structure of Pulmonary Embolism Response Teams associated with rates of advanced interventions and mortality? A multicenter retrospective study

Principal Investigators: Samuel G. Rouleau (Critical Care Fellow, UC Davis) and Angela Jarman (emergency medicine, UC Davis)

Co-Investigators: Christian Sebat (Pulm/Crit, UC Davis), Hannah Kelson (medical student, UC Davis), Brandon Eller (Emergency Medicine resident, UC Davis), Anthony Weekes (Emergency Medicine, Atrium Health), Daniel Troha (Emergency Medicine, Atrium Health), Christopher Kabrhel (Emergency Medicine, MGH), **David R. Vinson** (Roseville/Sacramento)

Study Sites: UC Davis, Atrium Health, MGH

Funding: None

Summary: We are seeking to investigate how the structure of PERTs impacts rates of advanced interventions (systemic thrombolytics, catheter-directed interventions, surgical embolectomy, ECMO) and mortality. While PERTs have proliferated since their introduction in 2016, the structure of PERTs varies by institution. We hypothesize that structure does not have an impact on mortality but does impact rate of advanced interventions, especially catheter-directed interventions. To answer these questions, we are conducting a retrospective cohort study across 3 separate institutions with different PERT models.

Status: We presented two abstracts at the Society for Academic Emergency Medicine meeting in May. We will be submitting two mss this summer.

34. Using a cancer-modified Pulmonary Embolism Severity Index score expands outpatient eligibility

Principal Investigator: **David R. Vinson** (Roseville/Sacramento)

Co-investigators: Aidan R. Campbell, Grace V. Heringer, and Camille J Smith, KP CREST Premed Research Interns, and Javeria Khader, UCD EM resident

Study Sites: KPNC

Summary: Society guidelines recommend validated risk stratification tools, like the Pulmonary Embolism (PE) Severity Index (PESI), to identify low-risk patients who may be eligible for outpatient care. In the original PESI, active and inactive cancer are treated as equivalent risk factors. Adverse outcomes in PE patients, however, correlate more strongly with active cancer (e.g., metastatic) than inactive cancer (e.g., history of only local disease, excised 10 years ago, without recurrence). Accordingly, a few studies and clinical pathways have modified PESI (mod-

PESI) by assigning cancer points only for active disease, not just for a cancer history. The degree to which mod-PESI expands outpatient eligibility has not been explored.

This is a secondary analysis of a retrospective cohort study conducted across 21 U.S. community hospitals from 2013-2019. It included ambulatory adults diagnosed with PE in primary care to evaluate outpatient management (discharge home) both from the clinic and, for those referred, from the ED. We included only those with vital signs (VSs) in the clinic or ED, using worst values for scoring. Active cancer was metastatic disease or recent (<12 months) diagnosis, anticancer treatment or palliative care. We will compare the proportion of those who were low-risk (mod-PESI vs PESI) in both the clinic and in the ED. We will also report 30-day major bleed, recurrent venous thromboembolism (VTE), and mortality among reclassified patients as a safety metric.

Status: Our high school research intern presented an abstract in March at the Western Regional meeting of the Society for Academic Emergency Medicine. One of our post-bac research interns presented this at the national meeting in Atlanta in May. The brief report is undergoing peer review.

35. Improving management of ED patients with unexplained syncope: prospective validation of the Canadian Syncope Risk Score

Principal Investigators: **David R. Vinson** (Roseville/Sacramento) and Mary E. Reed (KP Division of Research)

Co-Investigators: **Dana R. Sax** (Oakland/Richmond), Howard Dinh and **Erik R. Hofmann** (South Sacramento), **Nachi Gupta** (Redwood City), Madeline J. Somers (Division of Research), Annie Ma, Tracy Nguyen, and Xinna Chen (UC Davis EM residency), Leyla Farshidpour (UCSF Fresno), Isaac Wolfkind (MS2, California Northstate College of Medicine)

Study Sites: KP Oakland, Richmond, Roseville, Sacramento, and South Sacramento

Funding: The Permanente Medical Group Delivery Science Research initiative

Summary: The Canadian Syncope Risk Score looks promising as an accurate means of risk stratifying emergency department patients with acute unexplained syncope. But it has not been validated in a diverse U.S. population. This prospective study combines the Risk Score with multispecialty treatment recommendations in a web-based clinical decision support system and test its discrimination and calibration among 5 KPNC EDs. If the tool performs well, we will expand its use across the region.

Status: We implemented our electronic decision-support tool in 5 EDs in early 2022. Enrollment is complete. We continue to collect outcome data. We presented an abstract describing CDS impact on physician decision-making at ACEP in October 2024 and presented interim tool performance metrics at UC Davis in April. We presented an abstract at the regional meeting of the American College of Physicians at Stanford in October. We will be presenting the abstract at the Society for

Academic Emergency Medicine meeting in May. We are working now on the analysis of the complete dataset.

36. Clinical decision support to Optimize Care of patients with Atrial Fibrillation or flutter in the Emergency department: protocol of a stepped-wedge cluster randomized pragmatic trial (O'CAFÉ trial)

Principal Investigators: **David R. Vinson** (Roseville/Sacramento) and Mary E. Reed (Division of Research)

Co-Investigators: E. Margaret Warton, Mary E. Reed, and Adina S. Rauchwerger (DOR), the incredible Site Leads of the KP CREST Network EDs, along with Alan Go (DOR) and Matthew D. Solomon (Cardiology, Oakland). Thanks also to our students for their work on a review of AF guidelines: Disha Bahl (St. George's University School of Medicine), and Leyla Farshidpour (UCSF-Fresno), and Jennifer Zhang (now Harvard MPH graduate)

Study Sites: KP Northern California

Funding: The Permanente Medical Group Delivery Science Research initiative

Summary: Atrial fibrillation/atrial flutter (AF) is a clinical and socioeconomic burden to the U.S. healthcare system and will only worsen with the accelerated aging of the KP membership and U.S. population. Our prior research has identified suboptimal rate, rhythm, and stroke prevention treatments across KP Northern California EDs, along with twofold inter-facility variation in hospitalization rates of ED AF patients. This study will evaluate the impact of a web-based clinical decision support tool to improve the ED management of patients with primary AF.

Status: The clinical decision support tool is in use across 16 of 21 EDs. Our methods paper was published in *Trials*. Our ms on stroke prevention was published in *JAMA Netw Open*. We are undertaking analysis on a rate reduction study. We also are working on a review of AF guidelines to see how they speak to the issues that face emergency physicians in the management of AF.

37. Identifying barriers and facilitators to the evidence-based evaluation and management of febrile infants in the ED

Principal Investigator: Lauren Westafer (Emergency medicine, UMass Chan Medical School, Baystate)

Co-investigators: Taylor Bates (Pediatric emergency medicine Fellow, UMass Baystate), Blake Spirko (Pediatric EM, UMass Baystate), Natalie Strokes (Emergency medicine, UMass Baystate), **David R. Vinson** (Roseville/Sacramento)

Summary: We will conduct individual semi-structured interviews with emergency physicians at purposefully selected community and academic EDs in the United States to identify factors that influence the initial workup/evaluation of febrile infants and to understand clinical decision-making processes. We will map barriers and facilitators to the adoption of evidence-based guidelines in patient care using the Consolidated Framework for Implementation Research (CFIR).

Status: We have completed physician interviews and are analyzing data.

[Similar qualitative studies by Westafer and Vinson.](#)

38. ED care of oncology patients from a new ED/oncology collaboration

38A. Characteristics of pain-related ED visits in patients with cancer receiving treatment within an integrated health system

Principal Investigator: Raymond Liu (medical oncology, San Francisco)

Co-investigators: Andy Avins (DOR and primary care), Jenny Wei (internal medicine residency, KP San Francisco), Prince Wang (student, KP School of Medicine, Pasadena), **David R. Vinson** (Roseville/Sacramento)

Study Sites: KP Northern California

Funding: KPNC Graduate Medical Education, Kaiser Foundation Hospitals

Summary: Adult patients with active cancer undergoing systemic therapy often have challenging symptoms from their treatment or cancer itself. One especially burdensome symptom is pain: up to 50% of patients undergoing active therapy have cancer pain. Frequently, cancer pain prompts patients to seek acute medical care in the ED for pain relief. However, some patients may be better served in a different care setting. In this retrospective cohort study, we analyze KPNC patients with active cancer who have received systemic therapy at an infusion center from 2017 to through 2019 who subsequently (<30d) visited the ED with a chief complaint of pain and were treated with opioids. We describe the study population and analyze pain characteristics prior to and at the time of the ED visit. We also examine associations between patient characteristics with the risk and outcomes of ED visits for pain. Results may help identify individuals receiving infusion center care who are at risk of short-term ED presentations for pain treatment and may help guide future studies on care gaps contributing to inadequate pain control among our patients with cancer.

Status: Data collection is underway.

38B. Preventing Emergency Department Presentations of Lung Cancer in an Integrated Health System (PREVENT ED Study)

Principal Investigators: Brandon Anderson (resident, KP SFO) and Raymond Liu (medical oncology, San Francisco)

Co-investigators: Lori Sakoda, Mary E Reed, and Madeline J Somers (DOR), Arun Dang, Tina Huang, Seema Pursnani, and Cynthia Triplett (Santa Clara), **David R. Vinson** (Roseville/Sacramento)

Study Sites: KP Northern California

Funding: TPMG DARE Targeted Analysis Program

Summary: Lung cancer is the main cause of cancer mortality in the United States, contributing to more deaths than breast cancer, prostate cancer, and colon cancer combined. Studies have shown that patients who present and have their lung cancer diagnosed in the ED generally have a lower socioeconomic status, more advanced cancer stage, and higher rates of 1-year mortality. Smoking cessation efforts, newer treatments, and screening for lung cancer can improve patient outcomes; however, it is unknown whether these advances have had an equitable impact on ED diagnosis of lung cancer. We aim to characterize the population who presents to the ED for lung cancer diagnosis to understand if improved screening or early symptom identification would impact lung cancer outcomes and equity of care.

Status: We have presented abstracts at various oncology meetings. Our first ms is undergoing peer review.

39. Sexual health e-visit as a tool to improve access to prevention and treatment of sexually transmitted infections

Principal Investigators: Amanda Thornton, Dana Clutter (Infectious Diseases); Jacek Skarbinski (Infectious Diseases and Division of Research)

Co-investigators: Jonathan Volk, MD; Michael Silverberg, PhD; Christian Lee-Rodriguez, MD; Mitchell Luu, MD; Christine Bruno, PharmD; **David R. Vinson** (Roseville/Sacramento); Anne Srisuro, MD; Joshua Nugent, PhD

Study Sites: KP Northern California

Funding: TPMG Delivery Science Research initiative

Summary: Despite the clinical and public health importance of testing and treatment of sexually transmitted infections (STIs), including HIV, gonorrhea, chlamydia, and syphilis, and HIV pre-exposure prophylaxis (PrEP) use among at-risk individuals, STI testing and PrEP uptake rates remain low. To reduce barriers and increase access to sexual health services, the sexual health e-visit was launched in KPNC in February 2022 and is now used by our members approximately 10,000 times per month. The e-visit can be accessed through an online portal (KP.org) and allows members to obtain information on STIs, request appropriate STI testing after answering questions about sexual exposures, and self-refer for PrEP initiation. However, it remains unknown whether the sexual health e-visit increases access to care, STI testing, and PrEP uptake among at-risk individuals more efficiently and effectively than traditional office, telephone or email interactions with primary care providers and/or reduces ED utilization sexual health services. This retrospective cohort study will evaluate the role of the sexual health e-visit in members' access to sexual health services. Results will inform future e-visit development within KPNC and the use of similar platforms in other health care systems.

Status: Data collection is complete. We have been presenting preliminary findings to various departments across KPNC as well as at national meetings. The first manuscript is undergoing peer

review: “E-visit use for sexually transmitted infection (STI) screening in a large integrated healthcare system.”

40. Influenza-associated outcomes for children in the 2024-25 season stratified by current and previous season vaccination status

Principal Investigator: **Tony Zitek** (Modesto/Manteca).

Co-investigators: Alexander Yu, Laith Al-Najjar, and R. Grant Mello (Residents, KP Modesto); Yun-Yi Hung, Lorlelei Lee-Haynes and Nicola P. Klein (KP Division of Research)

Funding: Kaiser Permanente Northern California Graduate Medical Education, Kaiser Foundation Hospitals.

Summary: We are conducting a retrospective cohort study of pediatric patients to determine whether influenza vaccination during the 2023–24 season protected against influenza-related outcomes during the 2024–25 season. The primary outcome will be medically attended influenza infection confirmed by polymerase chain reaction. Secondarily, we will assess influenza-associated ED visits and hospitalizations. Patients will be categorized into four groups: unvaccinated in both seasons, vaccinated only in 2023–24, vaccinated only in 2024–25, or vaccinated in both seasons. Groups will be compared using unadjusted analyses and multivariable regression.

Status: A preliminary data analysis has been completed. We presented an abstract at the Society for Academic Emergency Medicine meeting in May.

41. Incidence and characteristics of cervical spine fractures in young children presenting to community hospitals

Principal Investigator: **Tony Zitek** (Modesto/Manteca)

Co-investigators: Zaid Alazawi and Edward Durant (Modesto/Manteca)

Study Sites: KP Northern California

Funding: Kaiser Permanente Northern California Graduate Medical Education, Kaiser Foundation Hospitals

Summary: In 2024, the Pediatric Emergency Care Applied Research Network (PECARN) published a cervical spine injury prediction rule for injured children. While reasonable, their data came entirely from level 1 pediatric trauma centers. It is unclear if this rule is applicable to patients who present to community hospitals where these injuries may be exceedingly rare, and mechanisms of injury may differ. We are conducting a retrospective assessment of patients aged 0-7 years old who presented to a KPNC ED for a cervical spine injury between October 1, 2010, and December 31, 2024. We will determine the incidence of cervical spine injuries in this population. We will

summarize the clinical characteristics of these patients and compare presentations for patients aged <2 to those 2-7 years old.

Status: A brief report was recently submitted for peer review in an emergency medicine journal.

Recent Publications (Q1 2026)^d

TPMG (Northern CA)

Ballard DW, Warton EM, Skarbinski J, Siqueiros MH, Cholleti SM, Vinson DR, Mark DG, Durant EJ, DiLena DD, Reed ME. [The long tail of Long COVID-19: Broad and extensive increase in utilization within an integrated healthcare system](#). *Cureus*. 2026;18(4):e106676.

Lam J, Rahman H, Brown MD, **Stuempfig ND**. [Implementation of a Resident Pod Associated With Increased Patient Encounters and Critical Procedures for Emergency Medicine Residents in a Community ED](#). *Cureus*. 2026;18(1):e101018.

Lippert SC, Xu RH, Tucker LS, DiLena DD, Rauchwerger AR, Rabbani J, **Kene MV**. [Patterns of chemical restraint use for emergency department patients with acute mental health distress](#). *Ann Emerg Med*. 2026;88(1):26-35.

Mark DG, Alavi M, Nugent JR, Reed ME, for the Kaiser Permanente CREST Network Investigators. [Sodium Correction Rates and Associated Outcomes Among Patients With Severe Hyponatremia: A Retrospective Cohort Study](#). *Ann Intern Med*. 2026;179(3):330-339.

KP DOR Spotlight: [Faster sodium correction tied to improved outcomes in patients with low sodium levels](#)

Annals Podcast: Centor RM, **Mark DG**. [Annals On Call - Hyponatremia Correction: How Fast Is Too Fast?](#) *Ann Intern Med*. 2026;179(2):e2600503OC.

McCullum M. [Redesigning emergency department processes: a nurse-led, portal-based flow model to optimise resource use and patient care](#). *Emerg Nurse*. 2026 Feb 24. Online ahead of print.

Sax DR, Warton EM, **Mark DG**, Vitale-McDowell TJ, DiLena DD, Rauchwerger AS, Reed ME. [Association between emergency department undertriage or overtriage with timeliness of care and patient outcomes](#). *Ann Emerg Med*. 2026;87(6):723-734.

Sax DR, Do T, Alavi M, Reed ME, Vora N, Gilbert AL, Mannis TE, Dokey AT, Vora RA. [Fundoscopic Cameras Associated with Shorter Emergency Department Length of Stay for Patients with Vision Loss](#). *Acad Emerg Med*. 2026;33(3):e70267.

^d A more comprehensive list of publications from the KP CREST Network can be found online: <http://www.kpcrest.net/>

El-Sabrou H, Lee D, Shah R, Gilbert AL, Patel A, **Sax D**, Reed M, Alavi M, Vora N, Hwang C, Mannis T, Dokey A, Moore S, Vora RA. [Branch retinal artery occlusion time to presentation and diagnosis: a retrospective review](#). *Int J Retina Vitreous*. 2026;12(1):49.

Lee D, El-Sabrou H, Shah R, Gilbert AL, Patel A, **Sax D**, Reed M, Alavi M, Vora N, Hwang C, Mannis T, Dokey A, Vora RA. [Branch Retinal Artery Occlusion Visual Outcomes: A Community-Based Study](#). *Ophthalmol Retina*. 2026;10(6):682-684.

Secko M, Singer DD, **Tetteh G**, Vitale M, Fombonne B, Bianconi K, Fernandes R, Gowie A, Singer AJ. [A Pilot Study Evaluating Erector Spinae Block Versus Saline For Emergency Department Patients With Ureterolithiasis](#). *J Emerg Med*. 2026;83:1-7.

Vinson DR, Rouleau SR, Balasubramanian MJ, Roubinian NH, Pai AP. [Weak evidence, rare eligibility, and complex care coordination limit application of structured surveillance for subsegmental pulmonary embolism](#). *Ann Emerg Med*. 2026;87(3):396-397.

Atzema CL, Cox JL, Cheung CC, Coll-Vinent B, Benjamin EJ, Jackevicius CA, **Vinson DR**. [Comparing guidelines for atrial fibrillation: focus on emergency medicine](#). *Ann Emerg Med*. 2026;87(4):435-450

Talerico R, de Wit K, Barco S, Bonorino J, den Uil C, Kraemer CE, Germini F, Iding A, Jones A, Konstantinides S, Masias C, Parks AL, Robert-Ebadi H, Tritschler T, Vedovati MC, **Vinson DR**, Woller SC, Klok FA. [Evidence-based risk stratification of patients with acute pulmonary embolism: Communication from the ISTH SSC Subcommittee on Predictive and Diagnostic Variables in Thrombotic Disease](#). *J Thromb Haemost*. 2026;24(3):1181-1189.

SCPMG (Southern CA)

Kerber KA, Sangha N, Burke JF, **Jancis MO**, Baecker A, Shen E, Nguyen H, Monjazebe S, Manthena P, Park S, Sharp AL, Meurer WJ. [Cumulative Incidence of Stroke Disability and Mortality Following Emergency Department Discharge for Dizziness: A Cohort Study](#). *Ann Emerg Med*. 2026;87(4):463-473.

Rose G, Shank W, **Esener D**. [Ultrasound-guided multifidus injection with 5 % dextrose for lumbar facet syndrome in the emergency department](#). *Am J Emerg Med*. 2026;99:518-520